

Letter to the Editor

Immune Cells Express Endocrine Markers

Michel Salzet

Received: January 13, 2002

Laboratoire de Neuroimmunologie des Annélides, UMR CNRS 8017, SN3, IFR 17 INSERM, Université des Sciences et Technologies de Lille, 59655 Villeneuve d'Ascq, France. TEL +33 3 2043 7277, FAX +33 320041130
E-MAIL michel.salzet@univ-lille1.fr

Abstract: Evidence are now given that immune cells expressed endocrine markers like neuropeptides, biogenic amine, neuropeptide processing enzymes, regulated secretion pathway. In clear, immune cells expressed like the nervous system an endocrine phenotype. This give the following question: what can we now consider as immune or endocrine?

Key words: **endocrinology; neuroimmunity; immunity**

As suggested by Kvetnoy and colleagues [1] in the letter to NEL editor entitled “Claude Bernard was right: hormones may be produced by « non-endocrine » cells”, Claude Bernard was the first in 1855 to suppose that not only endocrine glands but many organs in the organism have the ability for “internal secretion” [2]. The evidences that identical biogenic amines and peptides hormones are found in neurones and in amine precursor and uptake decarboxylation (APUD) cells located in different organs firstly confirmed his visions of the endocrinology [2]. The existence of a diffuse neuroendocrine system (DNES) concept was born [2]. Recently, Weigent and Blalock demonstrated that communication and reciprocal regulation between the nervous, endocrine and immune systems are essential for the stability of the organism, these three systems use the same signalling molecules. [3] Among others, cytokines, hormones and neuropeptides have been identified as messengers mediating the communication between the three systems [4–8]. Moreover, in the last few years various animal models have served to study neuroimmune mechanisms confirming the view of communication between the neuroendocrine and immune systems via neuropeptide signalling and through specific receptors [6–8].

Immune cells are able to synthesise neuropeptides acting as signalling molecules but also in organism defence [8–14]. Peptides with antibacterial properties have been shown to be derived from neuropeptide precursors such as proenkephalin and chromogranin B [8–14]. The role of neuropeptide precursors in immunity, through the release of antibacterial peptides, is an entirely novel concept. The biosynthetic pathway that leads to the production of biologically active neuropeptides begins with the synthesis of large inactive precursor proteins which are cleaved at specific paired or single basic residues within the Golgi secretory pathway [15]. It is a family of subtilase-like pro-protein convertases (SPCs) [15] that is largely responsible for these processing events that activate precursor proteins into neuropeptides. The SPCs

have been extensively studied in both neural and endocrine systems. However, much less is known concerning their expression, regulation and role within the immune system at the basal level [15, 16] or their function during microbial challenge [4]. We recently demonstrate that SPC functions are important since differential expression of SPCs and the resulting cleavage patterns determine the nature and biological activity of the peptide products. Thus, depending on the pattern of SPC expression, a single protein precursor can give rise to different peptides with diverse biological activities like antimicrobial substances (secretolytin, enkelytin/peptide B, dermaseptins [9–13, 17] or chemoattractant factors (Methionine-enkephalin) [7, 8].

Finally, the important point is the presence of at least four different types of granules formed during maturation of neutrophil granulocytes in the bone marrow [18]. Recent data support the concept that the (lysosomal type) azurophil granules are secreted through the endosomal pathway [19]. Taken together, these data suggest that regulated exocytosis from neutrophil granulocytes belongs to the general mechanism of secretion.

Considering all the above data, the aim of this letter is to point out that some revisions of the fundamental postulates of classical endocrinology need to be now undertaken.

Acknowledgements

The Centre National de la Recherche Scientifique, the MNERT supported this work.

REFERENCES

- Kvetnoy IM, Reiter RJ, Khavinson VK. Claude Bernard was right: hormones may be produced by « non-endocrine » cells. *Neuroendocrinol Lett* 2000; **21**:173–174.
- Bernard, C. *Leçons de physiologie expérimentale appliquée à la médecine*. Paris, 1855.
- Weigent DA, Blalock E. Production of peptide hormones and neurotransmitters by the immune system. *Neuroimmunoendocrinol*. 1997; **69**:1–30.
- Elmqvist JK, Scammel TE, Saper CB. Mechanisms of CNS response to systemic immune challenge: the febrile response. *Trends in Neurosci*. 1997; **20**:565–570.
- Salzet M, Vieau D, Day R. Cross-Talk Between Nervous and Immune Systems Through the Animal Kingdom: focus on opioids. *Trends in Neurosci*. 2000; **23**: 500–505.
- Salzet M, Breton C, Bisogno T, Di Marzo V. Comparative biology of endogenous Endocannabinoids system: possible role in immune response. *Eur. J. Biochem*. 2000; **267**:4917–4927.
- Stefano GB, Salzet M. Invertebrate opioid precursors: Evolutionary conservation and the significance of enzymatic processing. *Int Review of Cytol Endocrinol* 1999; **187**:261–286.
- Stefano GB, Cadet P, Scharrer B Stimulatory effects of opioid neuropeptides on locomotory activity and conformational changes in invertebrate and human immunocytes: Evidence for a subtype of delta receptor. *Proc Natl Acad Sci USA* 1989; **86**:6307–6311.
- Stefano GB, Salzet B, Fricchione G Enkelytin and opioid peptide association in invertebrate and vertebrate: Immune activation and pain. *Immunol Today* 1998; **19**:243–248.
- Goumon Y, Lugardon K, Kieffer B, Lefevre JF, Van Dorselaer A, Aunis D, Metz-Boutigue MH. Characterization of antibacterial COOH-terminal proenkephalin-A-derived peptides (PEAP) in infectious fluids. Importance of enkelytin, the antibacterial PEAP209-237 secreted by stimulated chromaffin cells. *J Biol Chem* 1998; **273**:29847–29856.
- Goumon Y, Strub JM, Moniatte M, Nullans G, Poteur L, Hubert P, Van Dorselaer A, Aunis D, Metz-Boutigue MH. The C-terminal proenkephalin-A diphosphorylated peptide (209–237) from adrenal medullary chromaffin granules possesses antibacterial activity. *Eur J Biochem* 1996; **235**:516–525.
- Tasiemki A, Salzet M, Benson H, Fricchione GL, Bilfinger TV, Aunis D, Metz-Boutigue MH, Goumon Y, Stefano GB. The Presence of Antibacterial Peptides in Human Plasma During Coronary Artery Bypass Surgery. *J Neuroimmunol*. 2000; **109**:228–235.
- Tasiemski A, Verger-Bocquet M, Cadet M, Goumon Y, Metz-Boutigue MH, Aunis D, Stefano GB, Salzet M. Proenkephalin and innate immunity in invertebrates: the antibacterial peptide, peptide B. *Mol. Brain Res*. 2000; **76**:237–252.
- Tasiemski A, Hammad H, Vandenbulcke F, Breton C, Bilfinger TV, Salzet M. Presence of chromogranin antimicrobial peptides in human plasma during coronary artery bypass surgery and evidence of an immune origin of these peptides. *Blood* (in press).
- Bergeron F, Leduc R., Day R. Subtilase-like pro-protein convertases: from molecular specificity to therapeutic applications *J Mol Endocrinol* 2000; **24**:1–14.
- Vindrola O, Padros MR, Sterin-Prync A, Ase A, Finkielman S, Nahmod V. Proenkephalin system in human polymorphonuclear cells. Production and release of a novel 1.0-kD peptide derived from synenkephalin. *J Clin Invest* 1990; **86**:531–537.
- Vouille V, Amiche M, Nicolas P. Structure of genes for dermaseptins B, antimicrobial peptides from frog skin. Exon-1 encoded prepropeptide is conserved in genes for peptides of highly different structures and activities. *FEBS Lett* 1997; **91**:27–32.
- Ligeti E, Mocsai A. Exocytosis of neutrophil granulocytes. *Biochem Pharmacol* 1999; **57**:1209–1214.
- Fittschen C, Henson PM. Linkage of azurophil granule secretion in neutrophils to chloride ion transport and endosomal transcytosis. *J Clin Invest* 1994; **93**:247–255.