

Innate Immunity in Lophotrochozoans: The Annelids

Michel Salzet^{1,*}, Aurélie Tasiemski¹, Edwin Cooper²

¹Laboratoire de NeuroImmunologie des Annelides, UMR CNRS 8017, SN3, Université des Sciences et Technologies de Lille, 59655 Villeneuve d'Ascq Cedex, France and ²Laboratory of Comparative NeuroImmunology, Department of Neurobiology, David Geffen School of Medicine at UCLA, University of California, Los Angeles, California 90095-1763, USA

Abstract: Innate immunity plays a major role as a first defense against microbes. Effectors of the innate response include pattern recognition receptors (PRR), phagocytic cells, proteolytic cascades and peptides/proteins with antimicrobial properties. Each element of these events has been well studied in vertebrates and in some invertebrates such as annelids. From these different researches, it appears that mammalian innate immunity could be considered as a mosaic of invertebrate immune responses. Annelids belonging to the lophotrochozoans' group are primitive coelomates that possess specially developed cellular immunity against pathogens including phagocytosis, encapsulation and spontaneous cytotoxicity of coelomocytes against allogenic or xenogenic cells. They have also developed an important humoral immunity that is based on antimicrobial, hemolytic and clotting properties of their body fluid. In the present review, we will emphasize the different immunodefense strategies that adaptation has taken during the course of evolution of two classes of annelids *i.e.* oligochaetes and achaetes.

INTRODUCTION

In the course of evolution annelids have developed different immunodefense strategies against microbes living in water or soil that are ingested during feeding or introduced into the body after injury [1]. For understanding immune processes, it is necessary to briefly review annelid anatomy. Earthworms (Oligochaeta, oligochaetes) and leeches (Hirudinea, achaetes) are characterized by the presence of two compartments containing free cells: (a) the blood system with haemocytes: this compartment does not seem to be actively implied in immunity and (b) the coelom that includes several coelomocytes populations playing a role in the immune defense (Fig. 1). We focus in the present review on Oligochaeta and Hirudinea in which most of the work on immunodefense has been performed. Both are important for certain biomedical applications [2-4].

I. CELLULAR IMMUNE RESPONSES IN ANNELIDS

A. Phagocytosis by LC and NK-like Lysis by SC

In oligochaetes, spontaneous allogeneic cytotoxicity of coelomocytes from *Eisenia fetida* and *Lumbricus terrestris* has been investigated *in vitro* using different assays (trypan blue, lactate dehydrogenase and ⁵¹Cr release; FACS analyses) [5, 6]. Suzuki and Cooper investigated xenogenic reactions against human tumor cells [6]. TEM and SEM studies showed that close contact of target cells with Lumbricus effector cells (coelomocytes with different mitochondrial properties) [7] is followed by lysis. The results confirm that effector cell/target cell contact is essential for cytotoxicity to

occur. These cytological events support analogies with cytotoxic activities that are affected when vertebrate natural killer cells are co-cultured with appropriate targets. Cytotoxicity supports the postulate of Cooper [8] and later Franceschi [9] that a primitive natural killer cell-like (analogue vertebrate NK cells) evolved early in phylogenesis [8-11] and the following results support this view. As in the NK system, the target specificity is exceedingly broad since xenogenic, allogenic, and even syngenic red blood targets are killed under appropriate conditions. The specificity of these cytotoxic cells may be directed at cell-surface glycoproteins on target cell surfaces since several defined mono- and disaccharides can block killing. Suzuki and Cooper examined whether modifying cell surface antigens can affect the level of cell-mediated cytotoxicity [12]. When co-cultured with small coelomocytes (SC) with NK-sensitive cells lines (K592), they become agitated, extending numerous pseudopodia that bind to and kill targets cells [13] (Fig. 2A, 2B). Large coelomocytes (LC) aggregate around the lysed targets and encapsulate them before forming granulomas [13]. These results suggest that phagocytosis and cytotoxicity are independently mediated by two different cell types; earthworm leukocytes as a group are polyfunctional and not exclusively phagocytic [4]. This supports the view of early divergence of phagocytosis and lysis.

The use of monoclonal antibodies directed against several human CD markers enabled Cossarizza *et al.* to focus on these two coelomocytes in *E. fetida* [14]: (1) the SC also called killer cells which are positive for CD11a⁺, CD45RA⁺, CD45RO⁺, CDw49b⁺, CD54⁺, 2-m⁺ and Thy-1⁺ (CD90) and (2) the phagocytic LC that are negative for these markers [15, 16]. Both cell types were negative for numerous other CD and MHC class I and class II markers. [17]. SC are active during recognition, rapidly binding to targets and LC are phagocytic. One cell group the SC effects recognition, bind-

*Address correspondence to this author at the Laboratoire de NeuroImmunologie des Annelides, UMR CNRS 8017, SN3, Université des Sciences et Technologies de Lille, 59655 Villeneuve d'Ascq Cedex, France; Tel: +33 3 2033 7277; Fax: +33 3 2043 4054; E-mail: michel.salzet@univ-lille1.fr




Fig. (1). Earthworm (*Lumbricus terrestris*) cross section through intestinal region. Annelids in general are characterized by a blood system (dorsal and ventral vessels) and a body fluid system contained in the coelomic cavities. Only the components of the coelomic fluid were shown to participate to the immune defense.

ing and killing, whereas cleaning up the remaining debris rests with the other cell type the LC.

In achaetes, differential CD staining at different time points following grafting experiments (four months between two transplants and specimen examination after 1 month), assume the presence of two different cell types, *i.e.* NK-like cells and CD8⁺ cells in the medicinal leech (generally classified as cytotoxic lymphocytes in vertebrates) [18, 19]. These coelomocytes are selectively stimulated by different antigens [19]. De Eguileor and colleagues using morphological and immunocytochemical approaches have analyzed the migratory behavior of cells involved in inflammatory responses of *Glossiphonia complanata* [18]. Like other annelids, leech leukocytes phagocytose and encapsulate foreign material selectively depending upon size [20]. Encapsulation is optimally visible when a parasite has been completely coated and isolated by a thick melanotic capsule [20]. Wounding in leeches is accompanied by "leucopoiesis", as it is generally accepted in vertebrates. In fact, in transplanted/wounded leeches, massive proliferation of lymphocyte-like cells stemming from a single cluster has been observed [20].

II. HUMORAL IMMUNE RESPONSES IN ANNELIDS

A. Is There Evidence of Complement Like Activity?

Both, cellular and humoral immune reactions in annelids are increased in response to an inflammatory or intracoelomic injection of foreign material. Certain annelid immune mechanisms, like NK-like activity and pore formation resemble functionally perforin and/or complement activity of vertebrates. The protective mechanisms of invertebrates consist, beyond cell immunity, of lysis and agglutination enhanced by components from the coelomic fluid like antigen binding protein, cytokines, antimicrobial substances.

Results in the field of research are numerous, especially in oligochaetes. In several experiments designed to test the possible existence of complement like activity, Cooper [21] co-cultured earthworm leukocytes with various NK dependent and independent targets [21]. Since washed coelomocytes were always used in cytotoxic assays, the experiments were done to test the capacity of cell-free supernatants. Adding supernatant from coelomocytes cultured for 10 min at 22°C caused significant cytotoxicity against K562 and other human cell lines. Cytotoxicity was not evident when supernatants from the same (autogenic) or different (allogenic) earthworms were added immediately after washing, nor did cytotoxicity occur when supernatants were heated at 56°C for 30 min before addition to K562 cultures. This suggests that a humoral factor-mediated component that may be complement like protein is involved in a significant percentage of target cell killings. Cytotoxicity cannot be interpreted as effected primarily by earthworm leukocytes.

B. Hemolytic Substances

1. A recognition Component

A coelomic cytolytic factor (CCF) has been isolated from coelomic fluid of the earthworm *E. fetida* [22]. CCF-1 shared functional analogy with vertebrate TNF based on N,N'-diacetylchitobiose lectin like domain/activity, despite a gene or protein homology [23]. This suggests that CCF and TNF lectin domain have been functionally conserved in the course of evolution as a recognition mechanism during innate immune responses [24]. CCF-1 is capable of lysing different mammalian tumor cell lines but by contrast with TNF, this cytolytic activity is mediated by lysis and is not due to proteolysis. Interestingly, CCF-1 recognizes lysozyme-predigested Gram-positive bacteria or the peptidoglycan constituent




Fig. (2). **A, B:** Earthworm cytotoxic cell activity; Small effector (E) coelomocyte in contact with K562 human targets (T), before (A) and after 5 min interaction (B). (with permission from [12]). **C:** Negative staining transmission electron microscopy of eiseniapore-treated sheep erythrocyte membranes. The image clearly shows many pore-like structures with an outer diameter of 10 nm. (with permission from [27]) **D.** Immune localization of theromacin and theromyzin, two antimicrobial peptides in the leech *Theromyzon tessulatum*. Theromacin and theromyzin signals are detected in large fat cells (LFC) in close contact with the body fluid of the leech. **E:** On semi-thin sections using an immunogold procedure, positive signals to theromacin were observed at the periphery of the LFC and in coelomic cavities (CC). **F:** Confocal microscopy analysis with anti-cystatin B (green) and anti-cathepsin L (red) in large coelomic cells of the leech *Theromyzon tessulatum*.

muramyl dipeptide as well as muramic acid (Fig. 3). The broad specificity of CCF for pathogen-associated molecular patterns results from the presence of two distinct pattern recognition domains. One domain, which shows homology with the polysaccharide and glucanase motifs of beta-1,3-glu-

canases and invertebrate defense molecules located in the central part of the CCF polypeptide chain, interacts with LPS and beta-1,3-glucans. The C-terminal tryptophan-rich domain mediates interactions of CCF with N,N'-diacetylchitobiose and muramic acid (Fig. 3). These data provide evidence for

the presence of spatially distinct carbohydrate recognition domains within this invertebrate immunodefense molecule [25].

2. Pore Forming Molecules

Besides CCF-1, pore-forming proteins have been characterized in the earthworm coelom: *i.e.* eiseniapore [26, 27], lysenin [28] and fetidin [29] (Table 1) These molecules bind to and disturb the lipid bilayer only when particular sphingolipids, (e.g. sphingomyelin), are present which are lacking in annelids [30, 31]. As for lysenin, cholesterol enhances eiseniapore lytic activity toward sphingomyelin-containing vesicles. Leakage of vesicles was most efficient when the lipid composition resembled that of the outer leaflet of human erythrocytes [30, 32]. Electron microscopy of erythrocyte membranes confirmed ring-shaped structures (pores). The channel complex consists of six monomers [26, 27] suggesting a plausible explanation for the mechanism by which components of the earthworm's immune system destroy non-self components (Fig. 3).

In addition, Kauschke *et al.* have demonstrated the existence of a perforin-like protein [33, 34]. Perforin and other lytic molecules seemed to be linked. One explanation is that perforin will generate the holes in target cell membranes facilitating the entry of the lytic molecules like lysenins, lysins, fetidins and the hemolysins (H₁, H₂, H₃)³³. [29, 35, 36] (Table 1). Each of these molecules has been demonstrated to be active towards certain human diseases (Niemann-Pick; blood clots).

C. Cytokines and Mitogenic Factors

Most of the data obtained at the present time on cytokines in invertebrates are by antibody recognition and not by sequence similarity. In annelids, the only data concerns the oligochaetes [37]. With respect to mitogenic factors, one was initially discovered in *Lumbricus* that exerts activity on T cells [38]. Later, a 60-kDa component with mitogenic activity on murine splenocytes was identified in the coelomic fluid of *E. fetida* and named CMF (Coelomic Mitogenic




Fig. (3). Hypothetical scheme of the putative innate immune response in earthworm involving cytotoxic factors. Septic injury provoke the release of Coelomocyte Cytolytic Factor I (CCF-I) from coelomocytes. CCF-I possesses two distinct domains *i.e.* the N-terminal domain interacts with lipopolysaccharide and 1,3-glucans and the C-terminal domain mediates its interactions with N, N-diacetylchitobiose and muranic acid [25]. CCF-I can trigger the prophenoloxdase cascade (pro-PO) implicating serine proteases and fetidins (Fe). CCF-I also enhances phagocytosis by its opsonizing properties and potentiates lysis against various erythrocytes. CCF-I can be considered as a pattern recognition molecule exerting a key role in the innate immune response of earthworm. Catalytic factors (fetidness, eiseniapore (EP), lysenins (Ly) are released from chloragocytes and coelomocytes [27-29]. These proteins bind to and disturb the lipid bilayer only when particular sphingolipids are present and perform pore channels.

Factor) [39]. CMF was found to bind ConA that could account for its ability to inhibit ConA-induced spleen cell proliferation. CMF is a trimer of a 20-kDa protein. N-terminal amino acid sequence of monomeric CMF reveals partial sequence homology with phospholipase A2 (PLA2). Moreover, CMF-enriched coelomic fluid exerts phospholipase activity comparable with that of bovine pancreatic PLA2. These results suggest that coelomic fluid of *E. foetida* contains an ubiquitous PLA2-like enzyme which might be involved in immune reactions in earthworms such as anti-bacterial mechanisms [39].

D. Antigen-Binding Proteins (ABP)

Laulan *et al.* reported synthesis by *L. terrestris* of specific substances in response to an immunization with a synthetic hapten [40]. Tuckova *et al.* identified a 56 kDa molecule, antigen-binding protein consisting of two disulfide-linked polypeptide chains (31 and 33 kDa) both of which participate in the formation of the antigen-binding site [41]. The kinetics of ABP formation revealed that the response reached a maximum level between the 4th and the 8th day after the first dose and approximately 4 days after the second challenge. The degree of specificity of the ABP after the secondary *in vivo* challenge increased, but even so it was considerably lower than that of vertebrate immunoglobulin [42].

E. Antimicrobial Proteins

The most studied antimicrobial protein in annelids is the lysozyme [43] (Table 1). This is an enzyme that cleaves the -1-4 bonds between N acetylglucosamine and N acetylmuramic acid of Gram positive bacterial cell walls. Lysozyme is often detected using its action against *Micrococcus lysodeikticus*.

In oligochaetes, background activity was 20-fold enhanced by one injection of either Gram positive or Gram negative bacteria and sheep red blood cells. Maximum induced activity occurred 4-5 hr after injection. Transcription and translation regulation of lysozyme have been investigated by Hirigoyenberry *et al.* [44]. Besides, lysozyme activity, the coelomic fluid of the earthworm *E. foetida andrei* exhibits antibacterial, hemolytic and hemagglutinating activities (See the cytotoxicity section). These activities are mainly mediated by two proteins, named fetidins, of apparent molecular mass 40 kDa and 45 kDa, respectively [29]. The recombinant protein was showed to inhibit *Bacillus megaterium* growth. Near its hemolysis activity, the protein also exhibits a peroxidase activity and a peroxidase signature was identified in residues 52-62 [29] (Table 1).

In achaetes, an intrinsic lysozyme-like activity was demonstrated for destabilase [45, 46]. Several isoforms of destabilase constitute a protein family with at least two members characterized by lysozyme activity [45, 46]. The corresponding gene family implies an ancient evolutionary history of the genes although the function(s) of various lysozymes in the leech remains unclear. Differences in primary structures of the destabilase family members and members of known lysozyme families allow one to assign the former to a new family of lysozymes. New proteins homologous to destabilase were recently described for *Caenorhabditis elegans* and

bivalve molluscs suggesting that the new lysozyme family can be widely distributed among invertebrates [45, 46]. Besides lysozyme, a 14 kDa protein sharing a bacteriostatic activity and belonging to the hemerythrin family has been found in fat cells [47] as well as glial cells of leeches (Ver-gote, unpublished data). Hemerythrin is a non-hemic-iron oxygen-transport protein acting as an iron scavenger towards bacteria in polychaetes [48] (Table 1).

F. Antimicrobial Peptides

During the last few years studies on the components of the innate immune system fully demonstrated the contribution of the antimicrobial peptides to host defense. Antibiotic peptides are small molecules (size comprised between 2kDa and 8kDa). Their implication in natural resistance to infection is sustained by their strategic location in phagocytes, in body fluids and at epithelial cell levels *i.e.* at interfaces between organisms and its environment [49] (Table 1).

In oligochaetes, an antimicrobial peptide named lumbricin I has been isolated and characterized from the earthworm, *Lumbricus rubellus*. Lumbricin I is a proline-rich antimicrobial peptide of 62 amino acids. Lumbricin I showed antimicrobial activity *in vitro* against a broad spectrum of microorganisms without hemolytic activity. A 29-amino acid peptide, named lumbricin I (6-34), which was derived from residues 6-34 of lumbricin I, showed marginally stronger antimicrobial activity than authentic lumbricin I. However lumbricin I gene is not induced by bacterial infection, but was constitutively expressed. Furthermore, the expression of lumbricin I gene was specific in adult *L. rubellus*, no expression was found in eggs and young *L. rubellus* [50].

In achaetes, Tasiemski *et al.*, have recently isolated four antimicrobial peptides from the coelomic fluid of the leech *Theromyzon tessulatum*: theromacin, theromyzin, peptide B [51-55] and lumbricin sharing high sequence homology with the one isolated from earthworm (unpublished data). Theromacin is a cationic 75 amino acids peptide with 10 cysteines residues arranged in a disulfide array showing no obvious similarities with other known antimicrobial peptides [53]. Theromyzin and peptide B are anionic peptides of 82 and 31 amino acid residues respectively [52, 53, 56]. Theromacin and theromyzin are processed from a larger precursor constituted by a signal peptide directly followed by the active peptide. Interestingly, peptide B is processed from the C terminal part of proenkephalin A (PEA), the major precursor of enkephalins. Peptide B and theromyzin possess bacteriostatic activity by contrast with theromacin which exhibit potent bactericidal activity towards Gram positive bacteria. Theromyzin constitutes the first anionic antimicrobial peptide evidenced in an invertebrate. Analysis of *theromacin* and *theromyzin* gene expression demonstrated both a developmental and a septic injury induction in large fat cells (LFC), which constitute a specific tissue that seems to be a functional equivalent of the insect fat body (Fig. 2D, 2E). Both peptides seem to exert their antimicrobial activity by a systemic action. All these data suggest for the first time an innate immune response in a lophotrochozoan comparable to the antibacterial response of an ecdysozoan *i.e.* *D. melanogaster*. Consequently, *Theromyzon* is an original invertebrate model which has developed two modes of fighting infections by antimicrobial peptides: (i) storage of antibacterial

peptide derived from PEA, particularly in coelomocytes and/or in nervous system, and release of the peptide into the coelomic fluid after immune challenge (ii) induction after septic injury of gene coding for more classical antimicrobial peptide, mainly in LFC, and rapid release into the body fluid of the antibiotic peptides (Fig. 4).

G. Cysteine and Serine Protease Inhibitors

Serine proteases, such as trypsin, thrombin, granzyme B, are important enzymes involved in regulating many physiologic events including the extrinsic pathways of blood coagulation and immune processes. In the immune system, killer lymphocytes combine the activity of a membranolytic protein, perforin, and the serine protease granzyme B to induce target cell apoptosis [57]. Similarly, cysteine proteases such as cathepsins are implicated in processing functions of mammalian antigen presenting cells (APCs) [58]. They allow Class II-antigenic peptide binding by degradation of Ii chaperone in the initial Class II-Ii complex [59].

In oligochaetes, three serine proteases have been isolated from *E. fetida* leukocytes and one is related to trypsin [60]. In the same species, a serine protease inhibitor of 14 kDa contained in the coelomic fluid has been characterized [61]. This enzyme inhibitor shares high homologies with several plant serine protease inhibitors. These molecules seem to be implicated in the cytotoxic cascade involving fetidins, CCF and the prophenoloxidase system (Fig. 3) [30].

In achaetes, a gene tandem cystatin B (*Tt-cysb*)-cathepsin L (*Tt-catl*) was identified in the leech *T. tessulatum* [62]. *Tt-cysb* belongs to the cysteine protease inhibitor family. *Tt-cysb* gene possesses an open reading frame of 306 nucleotides without signal peptide. The deduced sequence presents 54% sequence identity with human cystatin B. No homologs of *Tt-cysb* are present in *D. melanogaster* and *C. elegans* genomes. The active site sequence (QXVXG) is conserved as well as the glycine at position 5 allowing its interaction with its substrate, the cathepsin L. *Tt-cysb* constitutes the first cystatin B protein isolated in invertebrates. *Tt-catl* be-




Fig. (4). Hypothetical scheme of the putative innate immune response of leeches involving antibacterial peptides. Septic injury provokes an important production of mucous that would trap bacteria present in the external environment of the leech. Then, trapped bacteria would be killed by theromacin and theromyzin present in the mucous. At the same time, injury would induce theromacin and theromyzin gene expression and the secretion of the gene products from the LFC into the body fluid of the animal [53]. Thus, theromacin and theromyzin would exert their antimicrobial property through a systemic action. Moreover, antibacterial peptide (peptide B) as well as immune activator (Methionine Enkephalin (Met-Enk)) derived from proenkephalin A (PEA), particularly in coelomocytes and/or in nervous system, are released into the coelomic fluid after immune challenge [52]. This would lead, in conjunction with the phagocytic action of coelomocytes containing cystatin B (*Tt-cysb*) and cathepsin L (*Tt-catl*), to the microbial killing in the body fluid [62].

longs to the cysteine proteinase family and presents 68% sequence identity with *D. melanogaster* cathepsin L. *Tt-cysb* is only expressed in one circulating coelomic cells population. Bacterial challenge leads to an enhancement of the *Tt-cysb* transcript exclusively in these cells. Double *in situ* hybridization showed cathepsin L gene (*Tt-catl*) to be expressed in the same cells suggesting that both *Tt-cyst* and *Tt-catl* gene products might play a role in immune regulation through these circulating coelomic cells (Fig. 2F). These data are in line with those evidenced in mammals or in *Drosophila* in which cathepsin L is present in small granules of haemocytes and might play a role in phagocytic events [63].

III. PERSPECTIVES CONCERNING ANNELID IMMUNITY

Arthropods, especially insects, represent an important group of invertebrates because of their sheer numbers, and *Drosophila* has provided ample evidence that they possess effective innate immune responses. Yet it is unwise to over-generalize about the immune system of all invertebrates based upon this single species. If one looks at metazoan phylogeny based on small subunit rRNA, certain interesting revelations emerge. There are two major groups, the pre-laterians and the bilateria, the latter being subdivided into two major groups, the protostomes and deuterostomes. Of the protostomes, in turn, there are two subgroups, the Lophotrochozoa, to which the annelids and mollusks belong, and the Ecdysozoa to which the arthropods and nematodes belong. Although it will not be discussed here, the dominant body of evidence with respect to innate immunity and its relation to vertebrate (mammalian) and plants concerns findings in two ecdysozoan species, *Drosophila* and *C. elegans*. In contrast, lophotrochozoan systems show some distinct differences and may have solved problems of immunodefense in a special manner as has been revealed here with respect to two groups of annelids [3]. Annelids are among the first coelomates and are therefore of special phylogenetic interest. Polychaetes which we did not focus on in this review, are restricted to the marine domain and are considered as the most primitive annelids, based on morphology, physiology and development. Oligochaetes and achaetes are considered as being more evolved as reflected in their immune systems. Compared to other invertebrate groups, data on annelid's immunity reveal heavier emphasis on the cellular than on the humoral response suggesting that immune defense of annelids seems to be principally developed as cellular immunity [37]. However, the Expressed Sequence Tag (EST) sequencing programs on *Lumbricus terrestris* (LumbricBASE; <http://www.earthworms.org>); and the leech *Hirudo medicinalis* (<http://genome.uiowa.edu/projects/leech/>), combined with microarrays techniques developed on these models and imaging mass spectrometry will allow acquisition of more information about leech and earthworm humoral immunity. Our preliminary transcriptomic data obtained in leeches (Table 2) [62] dealing with immune responses suggest that this strategy is feasible and that substantial new knowledge will be obtained by analysing lophotrochozoans. This will only lead to further expansion of the two existing schemes that have been proposed concerning earthworm (Fig. 3) and leech (Fig. 4) immune responses. In this expansion, we predict more relevant biomedical/clinical applications further than what already exists.

ACKNOWLEDGEMENT

This work was in part supported by the CNRS, the MNERT, the FEDER, the Conseil Regional Nord Pas de Calais, the Genopole of Lille, the Plateau de Proteomique de l'USTL and the NIH-Fogarty Grant. E.L. Cooper is supported by the Alexander von Humboldt Foundation and GAAC grant from the Federal Republic of Germany and two grants from NATO, Cooperative Research Grant (971128) and Advanced Research Workshop (976680).

REFERENCES

References 64-66 are related articles recently published in Current Pharmaceutical Design.

- [1] Dhainaut A, Scaps P. Immune defense and biological responses induced by toxics in Annelida. *Can J Zool* 2001; 79: 233-253.
- [2] Sima P, Vetvicka V. Evolution of immune reactions. *Crit Rev Immunol* 1993; 13: 83-114.
- [3] Cooper EL, Kauschke E, Cossarizza A. Digging for innate immunity since Darwin and Metchnikoff. *Bioessays* 2002; 24: 319-33.
- [4] Cooper EL, Kauschke E, Cossarizza A. Annelid humoral immunity: cell lysis in earthworms. *Adv Exp Med Biol* 2001; 484: 169-83.
- [5] Valembais P, Roch, Boiledieu D. In Phylogeny of immunological memory, M.G. Manning, ed.; Elsevier-North Holland Biomedical Press: Amsterdam 1980; pp. 47-55.
- [6] Suzuki MM, Cooper EL. Allogeneic killing by earthworm effector cells. *Nat Immun* 1995; 14: 11-9.
- [7] Cossarizza A, Cooper EL, Quaglino D, Salvioli S, Kalachnikova G, Franceschi C. Mitochondrial mass and membrane potential in coelomocytes from the earthworm *Eisenia foetida*: studies with fluorescent probes in single intact cells. *Biochem Biophys Res Commun* 1995; 214: 503-10.
- [8] Cooper EL. Phylogeny of cytotoxicity. *Endeavor* 1981; 4.
- [9] Franceschi C, Cossarizza A, Monti D, Ottaviani E. Cytotoxicity and immunocyte markers in cells from the freshwater snail *Planorbis cornuus* (L.) (Gastropoda pulmonata): implications for the evolution of natural killer cells. *Eur J Immunol* 1991; 21: 489-93.
- [10] Cooper EL, Wright RK, Stein EA, Roch PJ, Mansour MH. Immunity in earthworms and tunicates with special reference to receptor origins, Karger 1987.
- [11] Cooper EL, Roch PG, Wright RK. Phylogeny of mononuclear phagocytes, University of Tokyo Press: Tokyo 1982.
- [12] Suzuki MM, Cooper EL. Spontaneous cytotoxic earthworm leukocytes kill K562 tumor cells. *Zoolog Sci* 1995; 12: 443-51.
- [13] Quaglino D, Cooper EL, Salvioli S, Capri M, Suzuki MM, Ronchetti IP, et al. Earthworm coelomocytes *in vitro*: cellular features and "granuloma" formation during cytotoxic activity against the mammalian tumor cell target K562. *Eur J Cell Biol* 1996; 70: 278-8.
- [14] Cossarizza A. Earthworm leukocytes that are not phagocytic and cross-react with several human epitopes can kill human tumor cell lines. *Exp Cell Res* 1996; 224: 174-182.
- [15] Roch P, Cooper EL, Eskinazi DP. Serological evidences for a membrane structure related to human beta 2-microglobulin expressed by certain earthworm leukocytes. *Eur J Immunol* 1983; 13: 1037-42.
- [16] Engelmann P, Pal J, Berki T, Cooper EL, Nemeth P. Earthworm leukocytes react with different mammalian antigen-specific monoclonal antibodies. *Zoology* 2002; 105: 257-265.
- [17] Cooper EL. In *Invertebrate immunology*, B and Muller Rinchevich, WEG, ed.; Springer-Verlag: Heidelberg 1996; pp. 10-45.
- [18] de Eguileor M, Grimaldi A, Tettamanti G, Valvassori R, Cooper EL, Lanzavecchia G. Lipopolysaccharide-dependent induction of leech leukocytes that cross-react with vertebrate cellular differentiation markers. *Tissue Cell* 2000; 32: 437-45.
- [19] de Eguileor M, Grimaldi A, Tettamanti G, Valvassori R, Cooper EL, Lanzavecchia G. Different types of response to foreign antigens by leech leukocytes. *Tissue Cell* 2000; 32: 40-8.
- [20] de Eguileor M; Tettamanti G; Grimaldi A; Congiu T; Ferrarese R; Perletti G, et al. Leeches: immune response, angiogenesis and biomedical applications. *Curr Pharm Des* 2003; 9: 133-47.

- [21] Cooper EL, Cossarizza A, Suzuki MM, Salvioli S, Capri M, Quaglino D, *et al.* Autogeneic but not allogeneic earthworm effector coelomocytes kill the mammalian tumor cell target K562. *Cell Immunol* 1995; 166: 113-22.
- [22] Bilej M, Brys L, Beschin A, Lucas R, Vercauteren E, Hanusova R, *et al.* Identification of a cytolytic protein in the coelomic fluid of *Eisenia foetida* earthworms. *Immunol Lett* 1995; 45: 123-8.
- [23] Beschin A, Bilej M, Brys L, Torreele E, Lucas R, Magez S, *et al.* Convergent evolution of cytokines. *Nature* 1999; 400: 627-8.
- [24] Beschin A, De Baetselier P, Bilej M. CCF, an invertebrate analogue of TNF, is not related to the other lytic components from *Eisenia foetida* earthworm. *Bioessays* 2002; 24: 974; author reply 975-6.
- [25] Bilej M, De Baetselier P, Van Dijk E, Stijlemans B, Colige A, Beschin A. Distinct carbohydrate recognition domains of an invertebrate defense molecule recognize Gram-negative and Gram-positive bacteria. *J Biol Chem* 2001; 276: 45840-7.
- [26] Lange S, Nussler F, Kauschke E, Lutsch G, Cooper EL, Herrmann A. Interaction of earthworm hemolysin with lipid membranes requires sphingolipids. *J Biol Chem* 1997; 272: 20884-92.
- [27] Lange S, Kauschke E, Mohrig W, Cooper EL. Biochemical characteristics of Eiseniapore, a pore-forming protein in the coelomic fluid of earthworms. *Eur J Biochem* 1999; 262: 547-56.
- [28] Sekizawa Y, Kubo T, Kobayashi H, Nakajima T, Natori S. Molecular cloning of cDNA for lysenin, a novel protein in the earthworm *Eisenia foetida* that causes contraction of rat vascular smooth muscle. *Gene* 1997; 191: 97-102.
- [29] Lassegues M, Milochau A, Doignon F, Du Pasquier L, Valembois P. Sequence and expression of an *Eisenia foetida*-derived cDNA clone that encodes the 40-kDa fetidin antibacterial protein. *Eur J Biochem* 1997; 246: 756-62.
- [30] Cooper EL, Roch P. Earthworm immunity: a model of immune competence. *Pedobiologia* 2003; 47: 676-688.
- [31] Milochau A, Lassegues M, Valembois P. Purification, characterization and activities of two hemolytic and antibacterial proteins from coelomic fluid of the annelid *Eisenia foetida*. *Biochim Biophys Acta* 1997; 1337: 123-32.
- [32] Shakor AB, Czurylo EA, Sobota A. Lysenin, a unique sphingomyelin-binding protein. *FEBS Lett* 2003; 542: 1-6.
- [33] Kauschke E, Pagliara P, Stanbili L, Cooper EL. Characterization of proteolytic activity in coelomic fluid of *Lumbricus terrestris* (Annelida, Lumbricidae). *Comp Biochem Physiol* 1997; 116: 235-242.
- [34] Porchet-Henneré EaD, T. Adaptability of the coccidian *Coelotropha* to parasitism. *Dev Comp Immunol* 1992; 16: 263-274.
- [35] Eue I, Kauschke E, Mohrig W, Cooper EL. Isolation and characterization of earthworm hemolysins and agglutinins. *Dev Comp Immunol* 1998; 22: 13-25.
- [36] Koenig S, Wagner F, Kauschke E, Peter-Katalinic J, Cooper EL, Eue I. Mass spectrometric analyses of CL(39), CL(41) and H(1), H(2), H(3) confirm identity with fetidin and lysenin produced by earthworm leukocytes. *Dev Comp Immunol* 2003; 27: 513-20.
- [37] Beck G, Habicht GS. Immunity and the invertebrates. *Sci Am* 1996; 275: 60-3.
- [38] Wojdani A, Stein EA, Alfred LJ, Cooper EL. Mitogenic effect of earthworm (*Lumbricus terrestris*) coelomic fluid on mouse and human lymphocytes. *Immunobiology* 1984; 166: 157-67.
- [39] Hanusova R, Bilej M, Brys L, De-Baetselier P, Beschin A. Identification of a coelomic mitogenic factor in *Eisenia foetida* earthworm. *Immunol Lett* 1999; 65: 203-11.
- [40] Laulan A, Lestage J, Chateaufreyaud-Duprat P, Fontaine M. [Substances contained in the coelomic fluid of *Lumbricus terrestris* having functions in common with those of some human complement components]. *Ann Immunol (Paris)* 1983; 134D: 223-32.
- [41] Tuckova L, Rejnek J, Bilej M, Pospisil R. Characterization of antigen-binding protein in earthworms *Lumbricus terrestris* and *Eisenia foetida*. *Dev Comp Immunol* 1991; 15: 263-8.
- [42] Bilej M, Tuckova L, Romanovsky A. Characterization of the limited specificity of antigen recognition in earthworms. *Folia Microbiol (Praha)* 1995; 40: 436-40.
- [43] Perin JP, Jolles P. The lysozyme from *Nephtys hombergi* (annelid). *Biochim Biophys Acta* 1972; 263: 683-9.
- [44] Hirigoyenberry F, Lassalle F, Lassegues M. Antibacterial activity of *Eisenia foetida* andrei coelomic fluid: transcription and translation regulation of lysozyme and proteins evidenced after bacterial infestation. *Comp Biochem Physiol B* 1990; 95: 71-5.
- [45] Baskova IP, Zavalova LL, Basanova AV, Sass AV. Separation of monomerizing and lysozyme activities of destabilase from medicinal leech salivary gland secretion. *Biochemistry (Mosc)* 2001; 66: 1368-73.
- [46] Baskova IP, Zavalova LL. Proteinase inhibitors from the medicinal leech *Hirudo medicinalis*. *Biochemistry (Mosc)* 2001; 66: 703-14.
- [47] Coutte L, Slomianny MC, Malecha J, Baert JL. Cloning and expression analysis of a cDNA that encodes a leech hemerythrin. *Biochim Biophys Acta* 2001; 1518: 282-6.
- [48] Deloffre L, Salzet B, Vieau D, Andries JC, Salzet M. Antibacterial properties of hemerythrin of the sand worm *Nereis diversicolor*. *Neuroendocrinol Lett* 2003; 24: 39-45.
- [49] Vizioli J, Salzet M. Antimicrobial peptides from animals: focus on invertebrates. *Trends Pharmacol Sci* 2002; 23: 494-6.
- [50] Cho JH, Park CB, Yoon YG, Kim SC. Lumbricin I, a novel proline-rich antimicrobial peptide from the earthworm: purification, cDNA cloning and molecular characterization. *Biochim Biophys Acta* 1998; 1408: 67-76.
- [51] Tasiemski A, Salzet M, Benson H, Fricchione GL, Bilfinger TV, Goumon Y, *et al.* The presence of antibacterial and opioid peptides in human plasma during coronary artery bypass surgery. *J Neuroimmunol* 2000; 109: 228-35.
- [52] Tasiemski A, Verger-Bocquet M, Cadet M, Goumon Y, Metz-Boutigue MH, Aunis D, *et al.* Proenkephalin A-derived peptides in invertebrate innate immune processes. *Mol Brain Res* 2000; 76: 237-52.
- [53] Tasiemski A, Vandenbulcke F, Mitta G, Lemoine J, Lefebvre C, Sautiere PE, *et al.* Molecular characterization of two novel antibacterial peptides inducible upon bacterial challenge in an annelid, the leech *Theromyzon tessellatum*. *J Biol Chem* 2004; 279: 30973-82.
- [54] Salzet M. Vertebrate innate immunity resembles a mosaic of invertebrate immune responses. *Trends Immunol* 2001; 22: 285-8.
- [55] Salzet M. Antimicrobial peptides are signaling molecules. *Trends Immunol* 2002; 23: 283-4.
- [56] Salzet M, Tasiemski A. Involvement of pro-enkephalin-derived peptides in immunity. *Dev Comp Immunol* 2001; 25: 177-85.
- [57] Trapani JA, Jans DA, Jans PJ, Smyth MJ, Browne KA, Sutton VR. Efficient nuclear targeting of granzyme B and the nuclear consequences of apoptosis induced by granzyme B and perforin are caspase-dependent, but cell death is caspase-independent. *J Biol Chem* 1998; 273: 27934-8.
- [58] Barrera C, Ye G, Espejo R, Gunasena S, Almanza R, Leary J, *et al.* Expression of cathepsins B, L, S, and D by gastric epithelial cells implicates them as antigen presenting cells in local immune responses. *Hum Immunol* 2001; 62: 1081-91.
- [59] Riese RJ, Chapman HA. Cathepsins and compartmentalization in antigen presentation. *Curr Opin Immunol* 2000; 12: 107-13.
- [60] Roch P, Cooper EL. Cellular but not humoral antibacterial activity of earthworms is inhibited by Aroclor 1254. *Ecotoxicol Environ Saf* 1991; 22: 283-90.
- [61] Roch P, Ville P, Cooper EL. Characterization of a 14 kDa plant-related serine protease inhibitor and regulation of cytotoxic activity in earthworm coelomic fluid. *Dev Comp Immunol* 1998; 22: 1-12.
- [62] Lefebvre C, Cocquerelle C, Vandenbulcke F, Hot D, Huot L, Lemoine Y, *et al.* Transcriptomic analysis in the leech *Theromyzon tessellatum*: involvement of cystatin B in innate immunity. *Biochem J* 2004; 380: 617-25.
- [63] Tryselius Y, Hultmark D. Cysteine proteinase 1 (CP1), a cathepsin L-like enzyme expressed in the *Drosophila melanogaster* haemocyte cell line mbn-2. *Insect Mol Biol* 1997; 6: 173-81.
- [64] Frantz S, Bauersachs J, Kelly RA. Innate immunity and the heart. *Curr Pharm Des* 2005; 11(10): 1279-90.
- [65] Furnkranz A, Leitinger N. Regulation of inflammatory responses by oxidized phospholipids: structure-function relationships. *Curr Pharm Des* 2004; 10(8): 915-21.
- [66] Hiemstra PS, Fernie-King BA, McMichael J, Lachmann PJ, Sallenave JM. Antimicrobial peptides: mediators of innate immunity as templates for the development of novel anti-infective and immune therapeutics. *Curr Pharm Des* 2004; 10(23): 2891-905.